

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07K 3/08, 15/06	A1	(11) International Publication Number: WO 92/10507 (43) International Publication Date: 25 June 1992 (25.06.92)
(21) International Application Number: PCT/US91/08971 (22) International Filing Date: 2 December 1991 (02.12.91) (30) Priority data: 624,169 7 December 1990 (07.12.90) US (71) Applicant: ABBOTT LABORATORIES [US/US]; CHAD 0377/AP6D-2, One Abbott Park Road, Abbott Park, IL 60064-3500 (US). (72) Inventors: BIENIARZ, Christopher ; 718 Old Trail, Highland Park, IL 60035 (US). HUSAIN, Mazhar ; 1643 Old Barn Circle, Libertyville, IL 60048 (US). BOND, Howard, E. ; 96 Little Melody Lane, Lake Forest, IL 60045 (US).		(74) Agents: GORMAN, Edward, H., Jr. et al.; Abbott Laboratories, CHAD 0377/AP6D-2, One Abbott Park Road, Abbott Park, IL 60064-3500 (US). (81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, KR, LU (European patent), MC (European patent), NL (European patent), SE (European patent). Published <i>With international search report.</i>
(54) Title: SITE-SPECIFIC CONJUGATION OF IMMUNOGLOBULINS AND DETECTABLE LABELS (57) Abstract <p>A process for producing a derivatized immunoglobulin and conjugates prepared therewith are provided. The derivatized immunoglobulin is produced by introducing sulfhydryl groups in the Fc region of an immunoglobulin wherein the Fab portion of the immunoglobulin is not altered. The derivatized immunoglobulin is used to prepare conjugates with various detectable moieties wherein a controlled distance exists between the Fc region of the immunoglobulin and the detectable moiety. The conjugates are particularly useful for performing immunoassays, and exhibit improved longevity, specificity and sensitivity.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU ⁺	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE*	Germany	MC	Monaco	US	United States of America
DK	Denmark				

⁺ Any designation of "SU" has effect in the Russian Federation. It is not yet known whether any such designation has effect in other States of the former Soviet Union.

- 1 -

SITE-SPECIFIC CONJUGATION OF IMMUNOGLOBULINS AND DETECTABLE LABELS

Field of the Invention

5

This invention relates to site-specific conjugation of immunoglobulins and detectable labels without alteration of the complementarity defined region of such immunoglobulins, and methods for preparing such conjugates. In particular, the present invention related to the conjugation of immunoglobulins and detectable labels at specific sites on the immunoglobulin while, at the same time, maintaining the accessibility of analyte binding sites on the immunoglobulin.

15

Background of the Invention

Although a variety of methods have been described for preparing conjugates of immunoglobulins and detectable labels, such as antibody-enzyme conjugates, such methods have resulted in random conjugation at many sites leading to heterogeneity and lack of reproducibility in the conjugates.

25

Site-specific carbohydrate labeling of the Fc region of antibodies with enzymes has also been described in O'Shannessy, et al., "Labeling of the Oligosaccharide Moieties of Immunoglobulins", Journal of Immunological Methods, Vol. 99 (1987) (pages 153-161).

30

However, conjugates prepared according to such method result in distances between the enzyme and the antibody which are intrinsically short, leading to distorted, conformationally strained antibodies and/or enzymes.

35

European Patent Application No. 32,863 describes conjugation methods and conjugation agents for

WO 92/10507

PCT/US91/08971

- 2 -

linking compounds containing carbohydrate moieties or
carboxyl groups to compounds containing thiol moieties
or electron-deficient moieties. In particular, the
coupling of immunoglobulins at the Fc region to labels,
5 such as enzymes, is described. However, such methods
and linking reagents used thereby provide reaction
conditions whereby there is essentially no
discrimination between the disulfides on the Fc region
and the complementarity defined region of the
10 immunoglobulin. This lack of discrimination results in
alteration of the complementarity defined region by
reduction of disulfide bonds therein to thereby affect
the binding sites on the immunoglobulin.


Accordingly, there is a need for a technique
15 to prepare site-specific conjugates which avoids
reductive cleavage of the disulfide bonds in, and
deterioration of, the complementarity defined region of
the antibody.

20 Summary of the Invention

According to the present invention, sulfhydryl
groups introduced in the Fc region of immunoglobulins
are coupled to enzymes wherein the Fab portion of the
25 immunoglobulin is not altered, and a controlled distance
exists between the Fc region and the enzyme label. As a
result, the conjugates prepared according to the present
invention exhibit greater binding capacity and signal
generating ability than those conjugates prepared
30 according to methods previously described. The
disadvantages generated by short, distorted and/or
conformationally strained conjugates are thus avoided,
while at the same time, improved conjugate longevity,
specificity, sensitivity, and other performance factors
35 result therefrom.

- 3 -

In particular, the sulfhydryl group is introduced through the Fc glycosylated portion of the immunoglobulin through a linking group containing a -NH-structure wherein the Fab portion, or complementarity defined region, is not altered. A presently preferred class of such substituted or derivatized immunoglobulin molecules is characterized by the formula:



wherein:

m is an integer from 1 through 20;

Y is an immunoglobulin molecule, wherein the Fc glycosylated region is reactive with a linker; and wherein either:

(i) X can be $-CH_2NH-$ or $-CH=NNHC-$;
 $\begin{array}{c} || \\ O \end{array}$

R₁ is a linking group containing 1 through 20 carbon atoms; and

n is an integer from 1 through 20; or

(ii) X-R₁-SH of formula (1) can be reduced glutathione or reduced hydrazidolipoamide.

The derivatized immunoglobulin molecule is readily coupled through its terminal sulfhydryl group to various detectable moieties, and the resulting conjugates are useful in a variety of diagnostic and therapeutic applications.

The detectable moieties which can be coupled to such a derivatized antibody (immunoglobulin) molecule are themselves first derivatized so as to contain, sulfhydryl-reactive functional groups which are preferably positioned as terminal groups, in a sterically unimpaired position, on the detectable moiety. Examples of such reactive functional groups


WO 92/10507

PCT/US91/08971

- 4 -

include haloacetates and preferably, maleimides. Techniques known in the art may be employed for preparing such derivatized moieties, and for preparing such conjugates. Preferably, conjugates according to the present invention comprise the derivatized molecules described above conjugated with an enzyme. The enzyme includes a coupling group, such as maleimide, haloacetyl (e.g., iodoacetyl, bromoacetyl and chloroacetyl), at the end of a long linking group. These conjugates are particularly useful in enzyme immunoassays.

Preferred conjugates contain a bifunctional coupling group of the structure:



wherein:

X and R_1 are as defined in Formula (1); and L is a linking group containing 1 through 40 carbon atoms, wherein L can be the same as or different than R_1 .

In Formula (2), X is joined to a moiety (Y) as defined in Formula (1) and L is joined to a detectable moiety (E).


More preferably, the class of such conjugates can be expressed by the formula:

35

WO 92/10507

PCT/US91/08971

- 5 -



5

wherein:

X is selected from the group consisting of

10 -CH₂NH-, -CH=NNHC-, a fragment of reduced glutathione,
and a fragment of reduced hydrazidolipoamide;

R₁ is a group containing 1 through 20
carbon atoms;


R₂ is cyclohexylmethyl;

15 R₃ is an alkylene group containing 1 through
10 carbon atoms; and

k is an integer of 1 through 10.

A presently preferred class of conjugates is
characterized by the formula:

20



25

wherein:

Q is the heterobifunctional coupling group of
Formula (3);

Y is as defined in Formula (1);

30

E is a detectable moiety;

n is as defined in Formula (1); and

o and p are each an independently selected
integer of 1 through 10.

35 Preferably, in Formula (4), the ratio of o to
p is in the range of from between about 1:1 to about

- 6 -

1:10, and n is in the range of from between about 1 to about 10. Conjugates of Formula (4) display improved homogeneity and lower polydispersity, and surprisingly improved sensitivity compared to conjugates prepared according to methods known in the art.

A derivatized immunoglobulin of the present invention is prepared by the sequential steps of site-specific periodate oxidation of immunoglobulin followed by reductive amination with an appropriate polyaminodisulfide reagent, and then disulfide reduction.

Brief Description of the Drawings

Fig. 1 illustrates the comparison of the activity of a carcinoembryonic antigen (CEA)/alkaline phosphatase conjugate prepared according to the present invention with an anti-CEA/alkaline phosphatase conjugate prepared according to a conjugation method known in the art.

Fig. 2 illustrates the comparison of the activity of an anti-cancer antigen (CA) 19-9/alkaline phosphatase conjugate prepared according to the present invention with an anti-CA 19-9/alkaline phosphatase conjugate prepared according to a conjugation method known in the art.

Fig. 3 illustrates the comparison of the activity of an anti-biotin/alkaline phosphatase conjugate prepared according to the present invention with an anti-biotin/alkaline phosphatase conjugate prepared according to a conjugation method known in the art.

Fig. 4 illustrates the comparison of the activity of a goat anti-human IgE/alkaline phosphatase conjugate prepared according to the present invention

- 7 -

with a goat anti-human IgE/alkaline phosphatase conjugate prepared according to a method known in the art.

Fig. 5 illustrates the comparison of the activity of an anti-pituitary thread protein (PTP)/alkaline phosphatase conjugate prepared according to the present invention with an anti-PTP/alkaline phosphatase conjugate prepared according to a conjugation method known in the art (values in parentheses represent enzyme to antibody ratios).

Description of the Invention

(a) Definitions

15

As used herein, the term "immunoglobulin" refers to a member of the class of protein molecules present, for example, in plasma, colostrum, tears, and other body fluids. An immunoglobulin, or antibody, combines specifically, non-covalently, and reversibly, with antigens.

20

The term "antibody" is used herein to refer to a homogeneous or heterogeneous population of immunoglobulins (such as monoclonal antibodies or polyclonal antiserum). An antibody-antigen complexation occurs by contact between specific combining sites of an immunoglobulin, known as complementarity defined regions, and antigenic determinants.

25

Five immunoglobulin classes are distinguished, identified as IgA, IgD, IgE, IgG and IgM. IgG is the predominant immunoglobulin in plasma, comprising about 75% of total plasma immunoglobulin, and is the most abundant immunoglobulin of internal body fluids, particularly extravascular materials, for combatting microorganisms and their toxins. There are four

30

35

- 8 -

subclasses of IgG, identified as IgG-1, IgG-2, IgG-3 and IgG-4, which are distinguished by differences in amino acid sequences. In human serum, IgG-1 comprises about 65% of plasma immunoglobulins.

5 As used herein, the term "Fc region" means the region of an immunoglobulin that corresponds to the stem or leg of the Y-shaped immunoglobulin molecule, and consists of the C-terminal sections of the two heavy chains linked by one or more disulfide bonds.

10 As used herein, the term "Fab region" means the region of an immunoglobulin other than the Fc region. Each Fab region consists of a light chain linked by a disulfide bond to the N-terminal part of a heavy chain, wherein each such Fab region is one of the
15 two arms of the Y-shaped immunoglobulin molecule. Each Fab region contains a single combining site in the highly variable region thereof, and behaves as a non-precipitating univalent antibody.

As used herein, the term "label" or
20 "detectable moiety" refers to a compound which can be used to label an immunoglobulin, usually by coupling such a compound with an immunoglobulin as in a conjugate. Labels can vary greatly in structure and function and include, but are not intended to be limited
25 to, enzymes, radiolabels, fluorogens, biotin, toxins, drugs, haptens, DNA, RNA, polysaccharides, polypeptides, liposomes, chromophores, chemiluminescers, colored particles and colored microparticles, and the like.

In the R_1 and R_2 group as defined above, for
30 example, in Formulas (1), (2), and (3), alkyl groups are preferred, cyclohexylmethyl being a particularly preferred R_2 group. The term "alkyl" includes straight and branched chains when three or more carbon atoms are present. Preferred alkyl chains contain less than seven
35 carbon atoms each. Presently most preferred are ethyl

WO 92/10507

PCT/US91/08971

- 9 -

and glutathione, wherein R_2 is preferably cyclohexylmethyl.

(b) Derivatized immunoglobulin

5

A derivatized immunoglobulin of this invention is prepared by the steps of:

- 10 (i) site-specific oxidation of the immunoglobulin to convert at least two hydroxyl groups in the glycosylated zone of the Fc region of the immunoglobulin to at least two aldehyde groups;
- (ii) reductive alkylation of the aldehyde groups to introduce disulfide moieties onto the Fc region of the immunoglobulin; and
- 15 (iii) reduction of the disulfide groups to sulfhydryl groups,

wherein aqueous liquid phase reaction conditions are employed in each step.

20 It is to be understood that oxidation of the immunoglobulin molecule can be accomplished by either chemical oxidation or enzymatic oxidation. For example, chemical oxidants include, but are not intended to be limited to periodate, bromine, and the like, and enzymatic oxidants include, but are not intended to be

25 limited to, the sequential use of neuraminidase and galactose oxidase, and the like.

Similarly, reduction of the disulfide groups of the immunoglobulin to generate terminal sulfhydryl groups can be accomplished by either chemically or

30 enzymatically cleaving the disulfide groups. For example, the chemical reduction of disulfide groups in the Fc introduced linkers may be accomplished by the use of mercaptoethanol, dithiothreitol (DTT), sodium borohydride, sodium dithionite, and the like. In the

35 case of enzymatic reduction of disulfide groups at the

WO 92/10507

PCT/US91/08971

- 10 -

5 Fc site, such reduction may be accomplished by exposing certain chemically introduced linkers containing the disulfide bonds to the appropriate enzymes which catalytically reduce the disulfides to thiols. Thus, the oxidized form of a linker-extended glutathione could generate thiols upon exposure to glutathione reductase. Similarly, the oxidized form of linker-extended lipoamide could generate thiols upon exposure to lipoamide dehydrogenase.

10 When the reductive alkylation is carried out with an amino disulfide compound, a Schiff base is formed which is stabilized by reduction.

After each of the indicated steps, the modified immunoglobulin product is preferably purified and concentrated according to methods known in the art.

15 According to the present invention, an alkali metal periodate, such as sodium periodate (NaIO_4), is employed to specifically oxidize vicinal diol moieties in a saccharide to aldehyde groups (see, for example, O'Shannessy et al.). The glycosylated zone of the Fc region of the immunoglobulin thereby undergoes a site-specific oxidation of at least two hydroxyl groups therein to two aldehyde groups. Accordingly, the complementarity defined region undergoes essentially no structural alteration. As will be understood by one skilled in the art, such oxidation is dependent on a number of variables including the concentrations of the immunoglobulin reactant and of the periodate oxidizing agent, pH, temperature and time. Various combinations of such variables may, of course, also be employed.

20 Some control of the extent of oxidation appears to be accomplished by varying the number of aldehyde groups generated and ultimately the degree of labeling of the oxidized immunoglobulin. The number of aldehyde groups generated for each immunoglobulin

25

30

35

WO 92/10507

PCT/US91/08971

- 11 -

molecule is dependent upon the oxidation conditions with more severe oxidation conditions generally producing more aldehyde groups per molecule. Preferably, the immunoglobulin contains from between about 2 to about 10
5 aldehyde groups for each oxidized immunoglobulin molecule in the glycosylated zone of the Fc region of the immunoglobulin.

The resulting aldehyde group-containing immunoglobulin is next most preferably purified and
10 concentrated. Purification is preferably accomplished by chromatography. For example, one presently preferred procedure is to pass the reaction medium containing the modified immunoglobulin through a gel filtration column such as a SEPHADEX[®] G-25 column (Pharmacia LKB
15 Biotechnology, Inc., Piscataway, NJ, USA) for purposes of separating the modified immunoglobulin. A column of such a composition is conventionally equilibrated and eluted using a buffer, such as sodium phosphate and sodium chloride having a pH of preferably about 7.0.
20 During elution, fractions are collected, and fractions having a suitable predetermined absorbance peak (e.g., 280 nm) are pooled. Protein concentration can be calculated from the absorbance peak using a suitable extinction coefficient.

25 Concentration of the aldehyde group-containing immunoglobulin from such a pool can be conveniently and preferably accomplished. A present preference is to employ centrifugation at about 5000 X g using tubes containing a molecular weight sizing membrane, such as
30 Centricon[®] tubes (Amicon Corp, Danvers, MA, USA) having the capacity to pass a material having a number average molecular weight cut-off of about 30,000.

Next, the purified, concentrated aldehyde group containing immunoglobulin is contacted under
35 aqueous liquid phase conditions with an appropriate

WO 92/10507

PCT/US91/08971

- 12 -

aminodisulfide reagent or a dihydrazido disulfide reagent to introduce at least one disulfide group for each immunoglobulin through the aldehyde groups.

Although it is possible to employ an amino or hydrazido substituted thiol (i.e. sulfhydryl) terminated compound, disulfide compounds are particularly preferred over thiol terminated compounds in order to maximize yields of the desired thiol-terminated group substituent, and to avoid undesired side reactions.

Accordingly, one preferred class of such disulfide compounds for use in reacting with an aldehyde group-containing immunoglobulin can be oxidized glutathione, hydrazidolipoamide, or a compound characterized by the formula:



where:

X is selected from the group consisting of $-CH_2NH_2$ and $-C(=O)-NHNH_2-$;

and

R_1 is a group containing 1 through 20 carbon atoms, preferably methylene.

As would be understood by one skilled in the art, the addition of bifunctional reagents to an ensemble of molecules causes both intramolecular modifications to the molecules and intermolecular modifications to the molecules. Accordingly, the aforementioned disulfide compounds of the present invention may similarly react with the carbohydrate-rich Fc regions of the immunoglobulin to result in either a large heterocyclic ring on the Fc region containing the disulfide bonds when reacting

WO 92/10507

PCT/US91/08971

- 13 -

intramolecularly, or may introduce a disulfide group between two immunoglobulins when reacting intermolecularly.

When the reaction with each aldehyde group on the immunoglobulin results in the formation of a Schiff base, the base may to be stabilized by reduction. For example, when reacted with a diamine disulfide reagent, stabilization by reduction of the Schiff base is required, whereas when the coupling reagent is dihydrazide disulfide, stabilization by reduction is not required. Such reductive stabilization is conveniently accomplished with an alkali metal cyanoborohydride, preferably sodium cyanoborohydride, under controlled reaction conditions. A present preference is to add the cyanoborohydride to the reaction medium within a time interval of from between about 5 minutes to about 24 hours after the addition of diaminodisulfide, more preferably from between about 15 minutes and about 60 minutes after such addition of diaminodisulfide. The reduction of the Schiff base results in the formation of a divalent secondary amine group.

The resulting derivatized disulfide group-containing immunoglobulin is preferably purified and concentrated according to the procedure and equipment described for use in the purification and concentration of the oxidized, aldehyde group-containing immunoglobulin.

Next, the purified, concentrated disulfide group-containing immunoglobulin is contacted under aqueous liquid phase conditions with a reducing agent which reduces the disulfide groups thereof to free sulfhydryl terminated groups. Where a chemical reducing agent is employed, a dithiol, such as dithiothreitol, is preferred. Where an enzymatic reducing reagent is employed, glutathione reductase, lipoamide

- 14 -

dehydrogenase, and the like, are preferred. The reduction of disulfide groups is achieved with essentially no effect upon other portions of the immunoglobulin molecule including the disulfide groups present in the internal structure thereof. This reduction is dependent on a number of variables, including the concentration of the dithiol reducing agent, as well as the pH, wherein combinations of disulfide reducing conditions can be employed. In this regard, the present inventors have surprisingly and unexpectedly found that at low enough concentrations of a reducing agent, such as dithiotreitol, (e.g., about 2 mM), only the disulfide bonds introduced by the present method into the Fc region undergo reduction, with the concomitant generation of thiols, wherein the disulfide bonds in the complementarity defined region and the hinge region are essentially unaffected at such low concentration of the reducing agent. Although an aldehyde group of the Fc region will react irreversibly with amine-containing linkers (e.g., cystamine and oxidized glutathione) only in the presence of sodium cyanoborohydride as a reducing reagent, sodium cyanoborohydride is not necessary when the linkers to be reacted with aldehydes of the Fc region contain a hydrazido group ($\text{H}_2\text{N}-\text{NH}-\overset{|}{\text{C}}=\text{O}$). Notably, use of sodium cyanoborohydride for reductive amination of the aldehydes on the Fc region does not reduce disulfide bonds in the reactive linkers or in the interchain links of the immunoglobulin.

According to the present invention, a substantially complete conversion of the Fc introduced disulfide side chain groups to sulfhydryl terminated side chains is accomplished. In contrast, reduction of the disulfide groups as previously described above (European Patent Application No. 32,863) occurs at

- 15 -

substantially higher concentrations of the chemical reducing agent.

5 The resulting sulfhydryl group containing immunoglobulin is preferably purified. A present preference is to utilize a purification procedure and equipment such as hereinabove described for use in the preparation of the oxidized, aldehyde group-containing immunoglobulin.

10 The purified product is directly suitable for conjugation with a detectable moiety.

(c) Conjugates of Derivatized Immunoglobulin With Detectable Moieties

15 A Fc site-specifically derivatized sulfhydryl group-containing immunoglobulin of the present invention can be conjugated with essentially any protein or other type of detectable moiety or label known in the art to produce new and useful conjugates. Such detectable
20 moieties or labels include, but are not intended to be limited to, enzymes, chromogens, luminescent, phosphorescent compounds, chemiluminescent compounds, fluorescent compounds, and the like. The sulfhydryl terminated side chain permits production of conjugates
25 wherein the unconjugated or normal activity of the starting immunoglobulin and the label are retained, and by employing the derivatized immunoglobulin of the present invention, conjugates are produced wherein constraints on conformational freedom are avoided so
30 that increased performance and stability can be achieved.

A label that is used for coupling with a derivatized immunoglobulin of the present invention is derivatized so as to contain, at least one terminal
35 sulfhydryl reactive functional group. Examples of such

- 16 -

reactive functional groups include, but are not intended to be limited to, maleimide, haloacetyl groups such as iodoacetyl, bromoacetyl and chloroacetyl, and the like. Techniques known in the art may be used for preparing such derivatized markers and for preparing such conjugates.

As described above, preferred conjugates of the present invention are those comprising an Fc site-specifically derivatized sulfhydryl group-containing immunoglobulin molecule coupled with a derivatized enzyme having a side chain coupling group that is terminated by a maleimide functional group. The coupling group employed in the enzyme is preferably chosen so as not to restrict the conformational freedom of the enzyme or its function after conjugating. In the coupling reaction, the terminal sulfhydryl group of the derivatized immunoglobulin molecule reacts with the terminal maleimide group of the derivatized enzyme to form the conjugate.

As would be understood by one skilled in the art, the present invention contemplates many different conjugates made with a coupling group as described herein. Many molecules of a single detectable moiety molecule can be attached to a single derivatized immunoglobulin molecule of this invention, or many derivatized immunoglobulin molecules can be attached to a single detectable moiety molecule. Also, a single conjugate of coupled molecules can contain a plurality of bifunctional coupling groups (Q) as described.


In preparing the preferred conjugates of the present invention, derivatized enzymes which have been made using the techniques described in copending U.S. patent application Serial No. 246,971, filed September 22, 1988, entitled "Heterobifunctional Coupling Agents", and incorporated herein by reference, can be used. Such

WO 92/10507

PCT/US91/08971

- 17 -

preferred starting enzymes are characterized by the formula:



wherein:

- 10 R_2 is a group containing 1 through 20 carbon atoms;
 R_3 is an alkylene group containing 1 through 10 carbon atoms;
 k is an integer from 1 through 10;
 15 n is an integer from 1 through 20; and
 E is an enzyme.

In the derivatized enzymes of Formula (7), the linking group between the enzyme (E) and the terminal maleimide group preferably contains a relatively long oligomeric chain of amino acids, from between about 2 and about 4 amino acids, which not only provides better physical separation between conjugated molecules incorporating such a linking group, but also functions to enhance the water solubility and dispersibility of a conjugate containing such chain.

20

25

To prepare a conjugate of a derivatized immunoglobulin of this invention and a derivatized enzyme having a side chain terminated by a maleimide group, any convenient procedure known in the art can be used. For example, a purified sulfhydryl group-containing immunoglobulin molecule is admixed under aqueous liquid phase conditions with a purified enzyme having a maleimide terminated side chain. Various combinations of reaction conditions may be employed in the coupling reaction of the sulfhydryl group with the

30

35

- 18 -

maleimide group. The coupling reaction is preferably carried out in the dark and is allowed to proceed for at least about 1 hour, preferably for about 12 hours. It is to be understood that periods of more than about 20
5 hours normally do not appear to increase the yield of conjugate. The conjugation or coupling reaction is believed to go substantially to completion based on starting molar ratios of reactant groups.

It is preferred to produce a conjugate having
10 no remaining free or reactive sulfhydryl groups. Accordingly, the unreacted sulfhydryl groups on the derivatized immunoglobulin molecules in the conjugate are reacted with a capping agent, such as maleimides substituted with alkyl groups of methyl through heptyl.
15 A preferred capping agent is N-ethyl maleimide (NEM). One convenient and presently preferred capping procedure is to add, to a buffered aqueous product conjugate-containing medium, an aqueous solution of NEM to a final concentration of about 0.3 mM, and agitate the mixture
20 for at least about one hour at ambient temperatures. The capped conjugate can be used directly for immunoassay purposes.

The conjugates according to the present invention are useful for the determination of an analyte
25 in homogeneous and heterogeneous immunoassay systems known in the art, such as competitive immunoassays, sandwich immunoassays, immunometric immunoassays, and the like, where the amount of a detectable moiety employed therein can be measured and correlated to the
30 amount of analyte present in a test sample. Generally, such measurement depends upon the ability of an immunoglobulin, i.e., an antibody, to bind to a specific analyte wherein a conjugate comprising an antibody to such analyte labeled with a detectable moiety is
35 employed to determine the extent of such binding.

- 19 -

Typically, the extent of binding is determined by the amount of the detectable moiety present in the conjugate which either has or has not participated in a binding reaction with the analyte, wherein the amount of the detectable moiety detected and measured can be correlated to the amount of analyte present in the test sample.

The conjugates according to the present invention are presented in a commercially packaged form as a composition or admixture where the compatibility of the reagents will allow or as a test kit, i.e., a packaged combination of one or more containers holding a conjugate according to the present invention and other appropriate reagents for performing an immunoassay as described herein and as known in the art. The reagents can further include other materials as are known to be useful in the assay from a user standpoint, such as buffers, diluents, standards, controls, and the like.

The present invention will now be illustrated, but is not intended to be limited, by the following examples:

Example 1

Site-specific Thiolation of Anti-Carcinoembryonic Antigen (Anti-CEA) Immunoglobulins and Its Coupling to Calf Intestinal Alkaline Phosphatase

(a) Introduction of Thiol Groups in the Fc region of the Antibody

A 1 mL solution of anti-CEA immunoglobulins (3-4 mg/mL in 0.1 M triethanolamine (TEA) buffer containing 0.16 M sodium chloride, pH 8) was placed in an amber vial. A 110 uL solution of 200 mM sodium-m-periodate (Sigma

- 20 -

Chemical Co., St. Louis, MO, USA) dissolved in 0.1 M TEA buffer, pH 8, containing 0.16 M NaCl was added to the antibody solution and the resulting mixture was gently stirred on a rotary agitator for 1 hour at approximately 5° C in the dark.

The oxidized antibody was purified from the above reaction mixture by chromatography on a 1 x 45 cm column of Sephadex G-25. The column was equilibrated and eluted with 0.1 M sodium phosphate, pH 7.0, containing 0.1 M NaCl. Fractions of about 1 mL each were collected during elution and the absorbance at 280 nm was determined. The peak fractions were pooled and the protein concentration of the immunoglobulins pool was calculated from its absorbance at 280 nm using an extinction coefficient ($E_{1\text{cm}}^{1\%}$) of 13.9.

The immunoglobulins pool was concentrated to 1.0 mL by centrifuging at 5000 x g using Centricon tubes (Amicon, Danvers, Massachusetts, USA) which contain a membrane sized to pass material having a number average molecular weight of up to about 30,000. The concentrate was admixed with 250 uL of 0.75 M cystamine dihydrochloride (Sigma Chemical Co.) dissolved in 0.1 M sodium phosphate, pH 7.0, containing 0.1 M NaCl, and the resulting mixture was gently stirred at room temperature. After about 15 minutes, 63 uL of 0.3 mM sodium cyanoborohydride (Sigma Chemical Co.) dissolved in 0.1 M sodium phosphate, pH 7.0, containing 0.1 M sodium chloride, was added and the resulting mixture was gently stirred on a rotary agitator overnight at room temperature.

The derivatized antibody was recovered by gel filtration on a 1 x 45 cm column of Sephadex G-25. The column was equilibrated and eluted with 0.1 M sodium phosphate, pH 7.0, containing 0.1 M NaCl and 2 mM EDTA (ethylene diamine tetraacetic acid). Fractions of about 1 mL each were collected during elution and the absorbance at 280 nm was determined. The peak fractions were pooled and

- 21 -

the concentration of the modified antibody was calculated from its absorbance at 280 nm as indicated above.

The pooled antibody was concentrated again, employing Centricon tubes of the same size and source as described above, to 1 mL and treated with 50 μ L of 40 mM dithiothreitol (Sigma Chemical Co.; dissolved in 0.1 M sodium phosphate, pH 7.0, containing 0.1 M NaCl and 2 mM EDTA) for 15 minutes at room temperature. The excess of dithiothreitol was removed by chromatography on a 1 x 45 cm Sephadex G-25 column. Fractions of about 1.0 mL each were collected and absorbance at 280 nm was measured. The peak fractions with absorbance at 280 nm of 0.3 or higher were pooled and stored on ice until the start of conjugation.

15 b) Derivatization of Enzyme

A 0.6 mL aliquot of calf intestinal alkaline phosphatase (Boehringer Mannheim, Indianapolis, Indiana, USA) at 10 mg/mL was diafiltered against 0.1 M sodium phosphate buffer, pH 7.0, containing 0.1 M NaCl, 1 mM $MgCl_2$, and 0.1 mM $ZnCl_2$. The volume was made up to 1 mL with the dialysis buffer and the enzyme solution was placed into a vial. To the enzyme solution was added 200 μ L of DMF containing 0.8 mg of succinimidyl (tricaproamido cyclohexylmethyl) N-maleimide (STCM) linker, a 30-atom extended heterobifunctional linker reagent with succinimide ester and maleimide termini, prepared as described in European Patent Application Publication No. 0314127 and incorporated herein by reference. The resulting reaction mixture was gently stirred on a rotary agitator for 30 minutes at room temperature and the derivatized enzyme was purified by chromatography on a 1 x 45 cm column of Sephadex G-25. The column was equilibrated and eluted with 0.1 M sodium phosphate buffer, pH 7.0, containing 0.1 M NaCl, 1 mM $MgCl_2$ and 0.1 mM $ZnCl_2$. Fractions of about 1 mL each were

- 22 -

collected during elution and the absorbance at 280 nm was determined. The peak fractions were pooled and the concentration of the enzyme in the pool was calculated from its absorbance at 280 nm using an extinction coefficient
5 ($E_{1\text{cm}}^{1\%}$) of 10.

c) Conjugation of Derivatized Enzyme with Antibody
Thiolated at the Fc region

10 The derivatized alkaline phosphatase from step (b) above was combined with the activated antibody from step (a) above in a molar ratio of 2:1 (enzyme to antibody). The mixture was gently stirred overnight at 2-8° C on a rotary agitator which resulted in approximately 90% conjugation of
15 the starting enzyme and antibody as indicated by HPLC and SDS-PAGE evaluations.

The unreacted thiol groups on the antibody were capped by treatment with N-ethylmaleimide (NEM; Sigma Chemical Co.) for a period of 1 hour at room temperature.
20 Aliquots of a 5 mM solution were added to the conjugate so that the final concentration of NEM in the conjugate was about 0.3 mM.

d) Performance of the Fc Conjugate

25 The performance of the Fc conjugate prepared as described above was evaluated by a sandwich-type microparticle capture enzyme immunoassay (MEIA) for carcinoembryonic antigen [CEA] (King, et al., In
30 Immunodiagnosis of Cancer, 2nd Ed., in press) using the Abbott IMx® System (Abbott Laboratories, Abbott Park, IL, U.S.A.; (Fiore, et al., Clin. Chem., Volume 34, pages 1726-1732, 1988). According to this assay format, latex microparticles covalently coupled with capture antibody
35 (anti-CEA) were incubated first with sample containing known

- 23 -

or unknown amounts of analyte (CEA) and then with alkaline phosphatase-labelled antibody conjugate (anti-CEA/alkaline phosphatase conjugate). Unadsorbed materials were removed at each step by capillary action and buffer washes.

- 5 Following the removal of unbound conjugate, enzyme substrate (4-methyl umbelliferyl phosphate) was added and rate of fluorescence increase was measured.

The data in Fig. 1 and Table 1 compare the performance of the Fc site-specific conjugate of the present invention with that of an anti-CEA/alkaline phosphatase conjugate prepared by combining iminothiolane-activated enzyme with STCM linker-activated antibody (STCM conjugate) as described in European Patent Application Publication No. 0314127. The results clearly demonstrate a 400% improvement in assay sensitivity with the Fc site-specific conjugate of the present invention over the STCM conjugate.

Table 1

20 Comparison of Anti-CEA/Alkaline Phosphatase Fc Site-Specific Conjugates with Conjugate prepared using Iminothiolane/STCM Extended Linker (STCM Conjugate) in IMx CEA Assay

25	CEA Standards (ng/mL)	Rate of Fluorescence (Counts/s/s)	
		STCM Conjugate (1 ug/mL)	Fc Conjugate (0.5 ug/mL)
30	0	9.0	19.1
	4	46.3	94.9
	10	97.4	200.6
	60	534.1	1000.3
	100	785.1	1565.7
35	200	1433.1	2634.3

- 24 -

The two conjugates described above were also each identically stressed for 3 days at 45° C to evaluate their stability. The data presented in Table 2 show that thermal stability of the Fc site-specific conjugate of the present invention was comparable to that of the STCM conjugate previously described, and that even after heat stress, the Fc site-specific conjugate of the present invention gave more than a 3-fold higher signal compared to the STCM conjugate.

Table 2

Comparison of Anti-CEA/Alkaline Phosphatase Fc Site-Specific Conjugate with STCM Conjugate after Heat Stress (3 Days at 45° C)

	CEA Standards (ng/mL)	Rate of Fluorescence (Counts/s/s)	
		STCM Conjugate (1 ug/mL)	Fc Conjugate (0.5 ug/mL)
0	0	12.7	26.1
4	4	37.6	66.2
10	10	79.2	132.6
60	60	377.9	616.3
100	100	599.1	968.3
200	200	1050.9	1669.6

- 25 -

Example 2

5 Fc Site-Specific Thiolation of Anti-Cancer Antigen (CA)
19-9 Immunoglobulins and Its Coupling To Calf Intestinal
Alkaline Phosphatase

10 An anti-CA 19-9 IgG and calf intestinal alkaline
phosphatase were each activated essentially as described in
steps (a) and (b), respectively, of Example 1, and then were
conjugated by combining them in a molar ratio of 1.5:1
(enzyme to antibody). The unreacted thiols were capped with
N-ethylmaleimide as described in Example 1.

15 The performance of the Fc site-specific conjugate was
analyzed using a sandwich-type MEIA format employing the
Abbott IMx system and compared to the performance of the
STCM conjugate prepared as described in Example 1. The
results presented in Fig. 2 and Table 3 demonstrate
20 approximately 8-fold higher sensitivity for the Fc site-
specific conjugate over the STCM conjugate.

WO 92/10507

PCT/US91/08971

- 26 -

Table 3

Comparison of Anti-CA 19-9/Alkaline Phosphatase Fc Site-Specific Conjugate with STCM Conjugates

5

	CA 19-9 Standards (Units/mL)	Rate of Fluorescence (Counts/s/s)	
		STCM Conjugate (4 ug/mL)	Fc Conjugate (1 ug/mL)
10			
	0	5.0	18.4
15	30	60.7	75.2
	90	141.1	237.8
20	180	276.0	462.7
	320	462.1	750.5
25	500	732.0	1072.2

Example 3

30

35 Fc Site-Specific Thiolation of Anti-Urinary Tract Infections (UTI) Immunoglobulins and Its Coupling To Calf Intestinal Alkaline Phosphatase

40 Rabbit and sheep anti-UTI immunoglobulins and calf intestinal alkaline phosphatase were activated essentially as described in steps (a) and (b), respectively, of Example 1, and then conjugated by combining the derivatized materials at a molar ratio of 2:1 (enzyme to antibody), and incubating overnight at 2-8° C with a gentle shaking on a rotary agitator. Following overnight incubation, residual

45

- 27 -

thiol groups in the conjugates were capped with 5 mM N-ethylmaleimide as described in Example 1.

The performance of such Fc conjugates was evaluated by Abbott Test-Pack™ immunoassay device [Abbott Laboratories, Abbott Park, IL, USA; W.E. Brown III, Clin. Chem., Volume 33, page 1567, (1987)] using E. Coli extract as the test organism and compared to that of a periodate conjugate prepared according to the method of Nakane and Kawaoi (P.K. Nakane and Akira Kawaoi, J. Histochem. and Cytochem. Volume 22, pages 1084-1091, 1974). The Fc site-specific conjugates were found to be about 4-fold more active than the periodate conjugate.

Example 4

15

20 Fc Site-Specific Thiolation of Anti-Biotin Immunoglobulins and Its Coupling To Calf Intestinal Alkaline Phosphatase

Rabbit anti-biotin immunoglobulins and calf intestinal alkaline phosphatase were activated essentially as described in steps (a) and (b), respectively, in Example 1, and then were conjugated by combining the derivatized materials in a 1:1 molar ratio. The resulting mixture was gently stirred overnight at 2-8° C on a rotary agitator. The following morning, unreacted thiols in the resulting conjugate were capped with 5 mM stock solution of NEM as described in Example 1.

The Fc site-specific conjugate when evaluated by DNA Probe Human Papilloma Virus (HPV) assay showed a 20-fold increase in sensitivity compared to the STCM conjugate prepared as indicated in Example 1 above (see Fig. 3 and Table 4).

WO 92/10507

PCT/US91/08971

- 28 -

Table 4

5 Comparison of Anti-Biotin/Alkaline Phosphatase Fc
 Site-Specific Conjugate with STCM Conjugates
 by HPV-DNA Probe Assay

Signal/Noise			
10	Molecules of HPV Target	STCM Conjugate (0.4 ug/mL)	Fc Conjugates (0.04 ug/mL) (0.2 ug/mL)
	0	1.0	1.0 1.0
15	100	1.0	1.08 1.17
	1000	1.0	1.40 1.92
20	10000	1.7	3.83 7.12

25

Example 5

30 Fc Site-Specific Thiolation of Anti-Cancer Antigen (CA 125)
 Immunoglobulins and Its Coupling To Calf Intestinal Alkaline
 Phosphatase

35 Monoclonal anti-CA 125 immunoglobulins and alkaline
 phosphatase were activated essentially as described in steps
 (a) and (b) respectively, of Example 1, and then were
 conjugated by combining the derivatized materials in
 respective molar ratios of 1:1 and 2:1 (enzyme to antibody).
 The resulting mixtures were gently stirred overnight at 2-8°
40 C on a rotary agitator. The resulting conjugates were then
 subjected to the end group capping procedure of Example 1 by
 adding 5 mM NEM.

 The performance of these conjugates was evaluated using
 a sandwich-type MEIA format employing the Abbott IMx System

- 29 -

and was found to be comparable to that of conjugates prepared by combining the DTT-reduced antibody with the enzyme derivatized with the STCM extended linker as described in European Patent Application Publication No. 0314127.

Example 6

10 Fc Site-Specific Thiolation of Anti-Human IgM
Immunoglobulins and Its Coupling To Calf Intestinal Alkaline Phosphatase

15 Goat anti-human IgM immunoglobulins and calf intestinal alkaline phosphatase were activated essentially as described in steps (a) and (b), respectively, of Example 1 and then were conjugated by combining the derivatized materials in
20 respective molar ratios of 1:1 and 2:1 (enzyme to antibody) and gently stirring the mixtures overnight at 2-8° C on a rotary agitator. The resulting conjugates were then subjected to the end group capping procedure of Example 1 by adding 5 mM NEM.

25 The performance of the Fc site-specific conjugates evaluated by a sandwich-type MEIA using the Abbott IMx system was found to be comparable to that of the conjugates prepared using iminothiolane and STCM extended linker as described in Example 1 above.

30

Example 7

35 Fc Site-Specific Thiolation of Anti-Human IgE
Immunoglobulins and Its Coupling To Calf Intestinal Alkaline Phosphatase

40 Goat anti-human IgE immunoglobulins and the alkaline phosphatase were activated essentially as described in steps

WO 92/10507

PCT/US91/08971

- 30 -

(a) and (b), respectively, of Example 1 and then were conjugated by combining the derivatized materials in respective molar ratio of 1:1. The antibody/enzyme mixture was gently stirred overnight at 2-8° C on a rotary agitator and the resulting conjugate was subjected to the end group capping procedure of Example 1 by adding 5 mM NEM.

The performance of the Fc conjugate was evaluated by a sandwich-type immunoassay as follows. A panel of allergens immobilized on a piece of nitrocellulose paper was treated with human IgE molecules specific to various allergens. The nitrocellulose paper was washed with phosphate buffer saline, treated with the Fc site-specific conjugates at various dilutions ranging from 1:1000 to 1:16000 following by color reaction for alkaline phosphatase. The results shown in Fig. 4 demonstrate that Fc site-specific conjugate is about 3-times more sensitive compared to the conjugates prepared by the periodate method of Nakane and Kawaoi referred to above.

20

Example 8

Site specific Thiolation of Anti-Pituitary Thread Protein (Anti-PTP) and Its Coupling To Calf Intestinal Alkaline Phosphatase

Anti-PTP immunoglobulins and the alkaline phosphatase were activated essentially as described in steps (a) and (b), respectively, of Example 1 and then were conjugated in a molar ratio of 1:1 and 2:1 (enzyme to antibody). The antibody/enzyme mixture was gently stirred overnight at 2-8° C on a rotary agitator and the resulting conjugate was subjected to the end group capping procedure of Example 1 by adding 5 mM NEM.

The performance of the Fc site-specific conjugate was evaluated using a sandwich-type MEIA format employing the

- 31 -

Abbott IMx System and compared to that of the periodate conjugate prepared according to the method of Nakane and Kawaoi referred to above. The results presented in Fig. 5 and Table 5 clearly show a 10-fold improvement in signal to noise ratio (S/N) of the Fc site-specific conjugate as compared to the periodate conjugate.

Table 5

10

Comparison of Anti-Pancreatic Thread Protein/Alkaline Phosphatase Fc versus Periodate Conjugate by IMx PTP Assay

15

PTP (ng)	Rate of Fluorescence (Counts/s/s)	
	Periodate Conjugate (1 ug)	Fc Conjugate (1 ug)
0.0	76.4	23.2
0.062	N.d.	66.3 (2.86)
0.125	N.d.	111.9 (4.82)
0.250	102.1 (1.34)	227.4 (9.80)
0.500	155.9 (2.04)	425.5 (18.34)
1.000	266.4 (3.49)	850.0 (36.64)

20

25

30

35

[The values in parentheses represent signal to noise ratios]

Example 9

40

Incorporation of Extended Analogs of Oxidized Glutathione into Fc region of Antibodies and Glutathione Reductase-Catalyzed Generation of Thiols

45

Antibodies are oxidized with sodium periodate at pH 8 and 2-8° C as described in Example 1 and then, after removal of excess reagent, subjected to reductive alkylation in the

- 32 -

presence of excess of oxidized glutathione (approximately 100 mM) and sodium cyanoborohydride (15 mM). After overnight incubation at room temperature, excess of reagent is removed by gel filtration over Sephadex G-25 as described in Example 1.

The final step in the activation of antibody is carried out enzymatically. To the derivatized antibody with extended analogs of oxidized glutathione attached covalently to the Fc region, are added a few micrograms of the enzyme, glutathione reductase and excess of NADPH (approximately 1 mM), and reaction is allowed to proceed until an adequate amount of thiols (2-6 moles/mole of antibody) are generated.

Subsequent to the generation of thiols, antibody is combined with alkaline phosphatase derivatized with STCM extended linker and the resulting mixture is gently stirred overnight at 2-8° C as described in Example 1. Finally the conjugate is capped with NEM as described in Example 1 and dialyzed to remove excess reagents.

Example 10

Incorporation of Hydrazido Lipoamide and Extended Analogs of Hydrazido Lipoamide into the Fc region of Antibodies and Lipoamide Dehydrogenase-Catalyzed Generation of Thiols

Antibodies are oxidized with sodium periodate at pH 8 and 2-8° C as described in Example 1, and then, after removal of excess periodate, treated with excess of hydrazido lipoamide or extended analogs of hydrazido lipoamide (10-50 mM). After overnight incubation at room temperature, excess of reagent is removed by gel filtration over Sephadex G-25 as described in Example 1.

The final step in the activation of antibody is carried out enzymatically. To the derivatized antibody with the lipoamide moiety covalently attached to the Fc region, are

- 33 -

added a few micrograms of lipamide dehydrogenase and excess of NADPH (approximately 1 mM) and reaction is allowed to proceed until an adequate number of thiols (2-6 moles of thiols/ mole of antibody) are generated.

5 Subsequent to the generation of thiols, antibody is combined with alkaline phosphatase derivatized with STCM extended linker and the resulting mixture is gently stirred overnight at 2-8° C as described in Example 1. Finally, conjugate is capped with NEM as described in Example 1 and
10 dialyzed to remove excess reagents before use.

 It will be apparent that many modifications and variations of the invention as herein set forth are possible without departing from the spirit and scope thereof, and that, accordingly, such limitations are imposed only as
15 indicated by the appended claims.

WO 92/10507

PCT/US91/08971

- 34 -

CLAIMS

We claim:


1. A process for producing a derivatized immunoglobulin, said process comprising the steps of:

5 (a) site-specifically oxidizing the glycosylated zone of the Fc region of an immunoglobulin with an oxidant to convert at least two hydroxyl groups in said zone of said immunoglobulin to an aldehyde group;

10 (b) reacting the oxidized immunoglobulin of step (a) with a disulfide compound wherein at least one of said aldehyde groups is reductively aminated to introduce a disulfide moiety onto the Fc region of said immunoglobulin; and

15 (c) selectively reducing the disulfide bonds of the disulfide moiety of the reacted immunoglobulin of step (b) with a reductant to convert at least one of said disulfide moieties of said immunoglobulin into two sulfhydryl groups.


2. The process of Claim 1 wherein said disulfide compound in step (b) is of the formula:



wherein

25 X is an aminoalkyl group or a semihydrazone group;
R₁ is a group containing from 1 through 20 carbon atoms; and R₁ and X together can form with the disulfide atoms between them a macrocyclic, heterocyclic ring or interimmunoglobulin disulfide containing chain to convert at least one aldehyde group per immunoglobulin molecule to a group which contains a

30 disulfide moiety of the formula:



wherein:

R₁ and X are as defined above.

- 35 -

3. The process of claim 1 wherein the oxidant in step (a) is a chemical oxidant.

5 4. The process of claim 2 wherein X is
selected from the group consisting of $\text{CH}_2\text{-NH}_2$, $\text{O}=\overset{\text{I}}{\text{CNHNNH}_2}$, a
fragment of glutathione, a fragment of reduced
hydrazidolipoamide, alkyl chain modified-length extended
10 glutathione, and alkyl chain modified-length extended
hydrazidolipoamide.

15 5. The process of claim 3 wherein the chemical oxidant
is selected from the group consisting of alkali metal
periodate, iodine and bromine.

6. The process of claim 3 wherein the chemical oxidant
is sodium periodate.

20 7. The process of claim 1 wherein the oxidant in step
(a) is an enzymatic oxidant.

25 8. The process of claim 7 wherein said enzymatic
oxidant comprises neuraminidase and galactose oxidase and
the oxidation step comprises sequentially contacting said
immunoglobulin with neuraminidase and galactose oxidase.

30 9. The process of claim 1 wherein the disulfide
compound in step (b) is selected from the group consisting
of cystamine, oxidized glutathione, oxidized hydrazido-
lipoic acid, alkyl chain modified-length extended oxidized
glutathione, and alkyl chain modified-length extended
oxidized hydrazidolipoic acid.

35 10. The process of claim 1 wherein the reductant in
step (c) is a chemical reductant selected from the group

- 36 -

consisting of mercaptoethanol, dithiothreitol, sodium borohydride and sodium dithionite.

11. The process of claim 1 wherein the reductant in
5 step (c) is an enzymatic reductant selected from the group consisting of glutathione reductase and lipoamide dehydrogenase.

12. The process of claim 1 wherein the sulfhydryl
10 group-containing immunoglobulin of step (c) is contacted with a detectable moiety containing a maleimide group coupled to said immunoglobulin to produce a conjugate.

13. The process of claim 12 wherein said detectable
15 moiety is selected from the group consisting of enzymes, chromophores, fluorescent molecules, chemiluminescent molecules, phosphorescent molecules, colored particles, and luminescent molecules.

14. The process of claim 12 wherein said detectable
20 moiety is an enzyme.

15. The process of claim 12 wherein said maleimide
25 group is joined to said detectable moiety through a linking group.

16. The process of claim 15 wherein said linking group is a heterobifunctional linking group.

17. The process of claim 1 wherein the reductant in
30 claim 1 is present in an amount whereby essentially no disulfide bonds of the native immunoglobulin in step (a) are reduced.

WO 92/10507


PCT/US91/08971

- 37 -

18. The process of claim 17 wherein the reductant comprises from between about 1 mM and about 5 mM dithiothreitol.

5 19. The process of claim 14 wherein the reductant comprises about 2mM dithiotreitol.

20. A derivatized immunoglobulin comprising an immunoglobulin substituted in the glycosylated zone of the Fc region thereof by at least one group of the structure:



wherein:


X is an aminoalkyl group or a semihydrazone group; and
R₁ is a linking group containing 1 through 20 carbon
15 atoms.

21. The derivatized immunoglobulin of Claim 20 wherein X is selected from the group consisting of -CH₂NH-,
-CH=NNHC-, alkyl chain modified-length reduced extended
20 $\begin{array}{c} \text{O} \\ \parallel \\ \text{O} \end{array}$
glutathione, and alkyl chain modified-length extended reduced hydrazidolipoamide.

22. The derivatized immunoglobulin of claim 20 wherein
25 R₁ is an alkyl group which contains 1 through 8 carbon atoms.

23. The derivatized immunoglobulin of claim 20 wherein
30 R₁ is methylene.

24. The derivatized immunoglobulin of claim 20 characterized by the formula:



35

WO 92/10507

PCT/US91/08971

- 38 -

wherein Y is an immunoglobulin molecule wherein the Fc glycosylated portion is reactive with a linking group and wherein:

- 5 m is an integer from 1 through 20; and
 n is an integer from 1 through 20.

10 25. A conjugate comprising the derivatized immunoglobulin of claim 20 coupled to a detectable moiety which contains a sulfhydryl-reactive functional group.


 26. The conjugate of claim 25 wherein said sulfhydryl-reactive functional group is maleimide.

15 27. The conjugate of claim 26 wherein said sulfhydryl-reactive functional group is linked to said immunoglobulin molecule by a bifunctional coupling group.

20 28. The conjugate of claim 25 wherein said detectable moiety is selected from the group consisting of enzymes, chromophores, fluorescent molecules, chemiluminescent molecules, phosphorescent molecules, colored particles and luminescent molecules.

25 29. The conjugate of Claim 25 wherein said detectable moiety is an enzyme.

 30. The conjugate of claim 22 wherein said bifunctional coupling group comprises the structure:




WO 92/10507

PCT/US91/08971

- 39 -

wherein X and R₁ are as defined in claim 20 and L, which can be the same as or different than R₁, is a linking group containing 1 through 40 carbon atoms.

5 31. A conjugate comprising an immunoglobulin coupled to a detectable moiety through at least one bifunctional coupling group having the structure:



15

wherein:


X can be -CH₂-NH- or -CH=NNC=O;

20 R₁ and R₂ are each an independently selected group containing 1 through 20 carbon atoms;

k is an integer from 1 through 10; and

X is joined to the glycosylated zone of the Fc region of said immunoglobulin, and the terminal >C=O group is
25 joined to said detectable moiety.

32. The conjugate of claim 31 characterized by the formula:



30 wherein:

Y is the glycosylated zone of the Fc region of an immunoglobulin molecule;

Q is said bifunctional coupling group;

E is an enzyme;

35 m is an integer from 1 through 20; and

WO 92/10507

PCT/US91/08971

- 40 -


o and p are each an independently selected integer from 1 through 10.

33. The conjugate of Claim 31 wherein said detectable moiety is selected from the group consisting of enzymes, chromophores, fluorescent molecules, chemiluminescent molecules, phosphorescent molecules, colored particles and luminescent molecules.

34. The conjugate of Claim 31 wherein said detectable moiety is an enzyme.

35. A method for the immunoassay determination of an analyte in a test sample, said method comprising the steps of:

(a) contacting said test sample with a conjugate comprising an immunoglobulin coupled to a detectable moiety through at least one bifunctional coupling group having the structure:



wherein:

X can be $-\text{CH}_2-\text{NH}-$ or $-\text{CH}=\text{NNHC}=\text{O}$;

R_1 and R_2 are each an independently selected group containing 1 through 20 carbon atoms;

k is an integer from 1 through 10; and


X is joined to the glycosylated zone of the Fc region of said immunoglobulin, and the terminal $>\text{C}=\text{O}$ group is joined to said detectable moiety;

- 41 -

(b) measuring the amount of said conjugate which either has or has not participated in a binding reaction with said analyte as a function of the amount of said analyte present in said test sample

5

36. The method claim 35 wherein said conjugate is characterized by the formula:



wherein:

10 Y is the glycosylated zone of the Fc region of an immunoglobulin molecule;

Q is said bifunctional coupling group;

E is an enzyme;


m is an integer from 1 through 20; and

15 o and p are each an independently selected integer from 1 through 10.

37. The method of claim 35 wherein said detectable moiety is selected from the group consisting of enzymes, 20 chromophores, fluorescent molecules, chemiluminescent molecules, phosphorescent molecules, colored particles and luminescent molecules.

38. The method of claim 35 wherein said detectable 25 moiety is an enzyme.

39. A test kit for the immunoassay determination of an analyte present in a liquid test sample, said test kit comprising a conjugate comprising an immunoglobulin coupled to a detectable moiety through at least one bifunctional 30 coupling group having the structure:



WO 92/10507

PCT/US91/08971

- 42 -

wherein:

5 X can be $-\text{CH}_2-\text{NH}-$ or $-\text{CH}=\text{NNHC}=\text{O}$


R_1 and R_2 are each an independently selected group containing 1 through 20 carbon atoms;

k is an integer from 1 through 10; and

10 X is joined to the glycosylated zone of the Fc region of said immunoglobulin, and the terminal $>\text{C}=\text{O}$ group is joined to said detectable moiety.

40. The test kit of Claim 39 wherein said conjugate is characterized by the formula:

15



wherein:

20 Y is the glycosylated zone of the Fc region of an immunoglobulin molecule;

Q is said bifunctional coupling group;

E is an enzyme;

m is an integer from 1 through 20; and

25 o and p are each an independently selected integer from 1 through 10.

41. The test kit of Claim 39 wherein said detectable moiety is selected from the group consisting of enzymes, chromophores, fluorescent molecules, chemiluminescent molecules, phosphorescent molecules, colored particles and luminescent molecules.

30

42. The test kit of Claim 39 wherein said detectable moiety is an enzyme.

35

1/5





Fig. 1

WO 92/10507

PCT/US91/08971

2 / 5



WO 92/10507

PCT/US91/08971

3/5




Fig. 3

4 / 5




Fig. 4

5/5




Fig. 5

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US91/08971

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ³		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC (5): C07K 3/08, 15/06 US CL : 530/390		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁴		
Classification System	Classification Symbols	
U.S.	424/85.91; 530/390	
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched ⁵		
Dialog Search, Biochem Database: S glutathione and (linker or spacer or linking) and aldehyde and reductase; s lipoamide and (linker or spacer or linking) and aldehyde		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ¹⁴		
Category ⁶	Citation of Document, ¹⁶ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No. ¹⁸
A	US, A, 4,140,679 (Malley) 20 February 1979.	4,9
A	US, A, 4,232,119 (Carlsson et al.) 04 November 1980.	1-42
Y	US, A, 4,699,784 (Shih et al.) 13 October 1987, see column 4, lines 7-14, column 6, lines 7-11.	5-8
X,P/ Y	US, A, 4,981,953 (Barbieri et al.) 01 January 1991, see column 1, lines 27-30, column 10, line 32- column 11, line 51, column 13, line 61- column 16, line 14, column 32, lines 4-8.	1 4,9,10,12,1515 -27,30/ 5-8,13,14,28, 29,31-42
Y,P	US, A, 4,994,385 (Bieniarz et al.) 19 February 1991, see the abstract, column 1, lines 29-39, Examples 7 and 8.	13,14,28, 29, 31-42
<p>* Special categories of cited documents:¹⁶</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search ²		Date of Mailing of the International Search Report ²
28 JANUARY 1992		26 MAR 1992
International Searching Authority ¹		Signature of Authorized Officer ²⁰
ISA/US		Jeffrey E. Russel 